



*The Alberta Lake Management
Society Volunteer Lake monitoring
report*

Wizard Lake

• • • • • • • • • •

2008 Report

Completed with support from:



Alberta Lake Management Society

Address: P.O. Box 4283
Edmonton, AB T6E4T3
Phone: 780-702-ALMS
E-mail: info@alms.ca

Water is integral to supporting and maintaining life on this planet as it moderates the climate, creates growth and shapes the living substance of all of Earth's creatures. It is the tide of life itself, the sacred source. David Suzuki (1997). The Sacred Balance.

Alberta Lake Management Society's Lakewatch Program

Lakewatch has several important objectives, one of which is to collect and interpret water quality data on Alberta Lakes. Equally important is educating lake users about their aquatic environment, encouraging public involvement in lake management, and facilitating cooperation and partnerships between government, industry, the scientific community and lake users. Lakewatch Reports are designed to summarize basic lake data in understandable terms for a lay audience and are not meant to be a complete synopsis of information about specific lakes. Additional information is available for many lakes that have been included in Lakewatch and readers requiring more information are encouraged to seek these sources.

ALMS would like to thank all who express interest in Alberta's aquatic environments and particularly those who have participated in the Lakewatch program. These people prove that ecological apathy can be overcome and give us hope that our water resources will not be the limiting factor in the health of our environment.

Acknowledgements

The Lakewatch program is made possible through the dedication of its volunteers and Lakewatch Chairs, Théo Charette and Ron Zurawell. We would like to thank Carole and LeVerne Ellsworth, Jerry McCracken, Matt Leach, and Bert Faulkner for their efforts in collecting data in 2008. We would also like to thank Lisa Brodziak and Sophie Damlencour who were summer interns with ALMS in 2008. Project Technical Coordinator, Jill Anderson was instrumental in planning and organizing the field program. Technologists, Shelley Manchur, Mike Bilyk, Brian Jackson and John Willis were involved in the logistics planning and training aspects of the program. Doreen LeClair and Chris Rickard were responsible for data management. Théo Charette (ALMS President) and Jill Anderson (Program Manager) were responsible for program administration and planning. Erika Brown, Zofia Taranu and Jesse Vermaire prepared this report, which was updated by Sarah Lord for the 2008 report. Alberta Environment and the Beaver River Watershed Alliance (BRWA) were major sponsors of the Lakewatch program.

Wizard Lake

Introduction

Wizard Lake is located in the North Saskatchewan River Basin, 50 km southwest of the city of Edmonton, in the counties of both Leduc and Wetaskiwin. Inspired by its meandering shape; the Indian name for this lake is “Lizard Lake” and was known as Conjuring Lake until the late 1960s. The year 1904 saw both the first settlers and the opening of a sawmill in the lake area. The sawmill was short-lived, closing in 1905, and succeeded by the opening of an underground coal mine, in operation until the 1940s. Today the area surrounding the lake includes Wizard Lake Jubilee Park and 110 cottages on the north shore, 61 cottages on the south shore, and a subdivision.




Figure 1. Bathymetric map of Wizard Lake. Each contour line represents a change in water depth of 1 meter.

Wizard Lake is a popular recreation spot for water skiing, canoeing, sailing, SCUBA diving, and fishing. Yellow perch (*Perca flavescens*) and northern pike (*Esox lucius*) are the most commonly fished species in the lake. There was a proposal to limit boat speeds to 12 km/hr in the west half of the lake in the late 1970's; however, whether this is enforced today is unclear (Mitchell & Prepas, 1990).

Wizard Lake occupies an area of 2.48 km², with a maximum depth of 11 m and mean depth of 6.2 m. The length of the lake stretches 11.5 km and has a maximum width of 0.55 km. It is a eutrophic lake, usually clear, but experiences dense blue-green algal blooms during the summer months, turning the water murky. The lake basin is made up of approximately 65% forest or bush, 25% agricultural land, 7% lake or sloughs, and 3% urban development (Mitchell & Prepas, 1990).

Results

Water Level

Water levels are measured as the elevation in meters above sea level (m asl) of the surface of the lake. Since the start of water level monitoring for Wizard Lake the water level has remained relatively constant (**Figure 2**). Fluctuations in the water level from year to year are rarely greater than 0.2 m and total change in water level over the entire monitoring record has been negligible.




Figure 2. Historical water levels (meters above sea level (asl)) in Wizard Lake, Alberta 1968 – 2009.

Water Temperature and Dissolved Oxygen

Water temperature and dissolved oxygen profiles in the water column can provide information on water quality and fish habitat. Please refer to the end of this report for descriptions of technical terms.

Temperature profiles taken for Wizard Lake throughout the summer varied little (**Figure 3**). The temperature of the water column in May and June gradually declined with increasing depth. Strong thermoclines have been observed on Wizard Lake in previous summers, but in 2008 a very weak thermocline was present only in late July at 5.5 m below the lake surface. The water column returned to a completely unstratified state in August and September.

In contrast, the dissolved oxygen concentrations in Wizard Lake were widely varying through the summer. Surface waters were well-oxygenated in late May and mid-June, but

a chemocline was present at 7.5 m depth and waters were anoxic below 8 m depth. In July, surface waters were highly oxygenated (oxygen is a byproduct of algal photosynthesis) and the chemocline had risen to 4.5 m depth, but lake waters were anoxic below 6 m depth. By mid-August, dissolved oxygen concentrations in surface waters had dropped below June levels, and the chemocline had dropped back to 7.5 m. Before 2 September a significant mixing event (fall turnover) occurred to equalize the oxygen concentrations in the top 10m of the water column. The top 5 m of Wizard Lake was well within surface water quality guidelines for Alberta for the duration of the summer (Figure 3).




Figure 3. Water temperature (°C) and dissolved oxygen (mg/L) profiles for Wizard Lake during the summer of 2008.

Water clarity and Secchi Depth

Water clarity is influenced by suspended materials, both living and dead, as well as dissolved coloured compounds in the water column. During the melting of snow and ice in spring, lake water can become turbid (cloudy) from silt transported into the lake. Lake water usually clears in late spring but then becomes more turbid with increased algal biomass as the summer progresses. The easiest and most widely used measure of lake water clarity is the Secchi disk depth.

During the 2008 open water season, the average Secchi disk depth of Wizard Lake was 1.43 m (Table 1). Maximum water clarity was observed in mid-June with a Secchi disk depth of 1.75 m. Water clarity declined to 1.25 m through mid-July and August, and returned to 1.3 m in September. The relatively constant water clarity through the summer indicates that constant mixing occurred throughout the open water season. Declines in water clarity in July and August were associated with a strong increase in algal biomass.

Water chemistry

Based on the trophic status of lake water characteristics, Wizard Lake is considered to be eutrophic to hyper-eutrophic (nutrient rich; see *A Brief Introduction to Limnology* at the end of this report). The concentrations of total phosphorus ranged from 39 to 59 $\mu\text{g/L}$ during the open water season, whereas chlorophyll *a* (a measure of water greenness and algal biomass) increased from 11.5 $\mu\text{g/L}$ in early June to 34.5 $\mu\text{g/L}$ by mid-September (**Figure 4**). Given that lakes in Alberta are productive, in the context of this province, Wizard Lake is about average in nutrient concentration and algal production.

Wizard Lake is well protected from acidification; its pH of 8.3 is well above that of pure water (i.e., pH 7; **Table 1**). Bicarbonate, calcium, magnesium, and sodium are the dominant ions in Wizard Lake. In general, there is a lack of water quality information for Wizard Lake to help examine long-term trends.




Figure 4. Total phosphorous, chlorophyll *a* (a measure of algal biomass), and total nitrogen concentrations for Wizard Lake during the summer of 2008.

Table 1. Mean values from summer 2006 samples, Wizard Lake.

Parameter	2006	2008
TP ($\mu\text{g/L}$)	48.4	50.2
TDP ($\mu\text{g/L}$)	13.6	12.4
Chl- <i>a</i> ($\mu\text{g/L}$)	32.6	23.9
Secchi disk depth (m)	1.33	1.43
TN (mg/L)	1300	1216
NO ₂₊₃ ($\mu\text{g/L}$)	7	<0.005
NH ₄ ($\mu\text{g/L}$)	31.4	20.6
Ca (mg/L)	25	27.9
Mg (mg/L)	8.5	8.9
Na (mg/L)	36	34.9
K (mg/L)	6	5.8
SO ₄ (mg/L)	3.5	4.5
Cl (mg/L)	4.7	4.5
CO ₃ (mg/L)	6	10
HCO ₃ (mg/L)	202	206.3
pH	8.3	8.3
Conductivity ($\mu\text{S/cm}$)	335	337.3
Total dissolved solids (mg/L)	186	191
Total Alkalinity (mg/L CaCO ₃)	172	175

Note: TP = total phosphorus, TDP = total dissolved phosphorus, Chl-*a* = chlorophyll *a*, NO₂₊₃ = nitrate+nitrite, NH₄ = ammonium, Ca = calcium, Mg = magnesium, Na = sodium, K = potassium, SO₄ = sulphate, Cl = chloride, CO₃ = carbonate, HCO₃ = bicarbonate.

*Atlas of Alberta Lakes (Mitchell and Prepas, 1990).

Works Cited

Mitchell, P. and E. Prepas. 1990. Atlas of Alberta Lakes. University of Alberta Press.

A brief introduction to Limnology

Indicators of water quality

The goal of **Lakewatch** is to collect water samples necessary to determine the water quality of lakes. Though not all encompassing, the variables measured in **Lakewatch** are sensitive to human activities in watersheds that may cause impacts to water quality. For example, nutrients such as phosphorus and nitrogen are important determinants of lake productivity. The concentrations of these nutrients in a lake are affected (typically elevated) by land use changes such as increased crop production or livestock grazing. Elevated nutrient concentrations can cause increases in undesirable algae blooms resulting in low dissolved oxygen concentrations, degraded fish habitat and production of noxious odors. Large increases in nutrients over time may also indicate sewage inputs, which in turn, may result in other human health concerns such as harmful bacteria or protozoans (e.g. *Cryptosporidium*).

Temperature and mixing

Water temperature in a lake dictates the behavior of many chemical parameters responsible for water quality (Figure 6). Heat is transferred to a lake at its surface and slowly moves downward depending on water circulation in the lake. Lakes with a large surface area or a small volume tend to have greater mixing due to wind. In deeper lakes, circulation is not strong enough to move warm water to depths typically greater than 4 or 5 m and as a result cooler denser water remains at the bottom of the lake. As the difference in temperature between warm surface and cold deeper water increases, two distinct layers are formed. Limnologists call these layers of water the **epilimnion** at the surface and the **hypolimnion** at the bottom. The layers are separated by a transition layer known as the **metalimnion** which contains the effective wall separating top and bottom waters called a **thermocline**. A thermocline typically occurs when water temperature changes by more than one degree within one-meter depth. The hypolimnion and epilimnion do not mix, nor do elements such as oxygen supplied at the surface move downward into the hypolimnion. In the fall, surface waters begin to cool and eventually reach the same temperature as hypolimnetic water. At this point the water mixes from top to bottom in what is called a **turnover** event. Surface water cools further as ice forms and again a thermocline develops this time with 4° C water at the bottom and 0° C water on the top.




Figure 6: Difference in the circulation of the water column depending on thermal

In spring another turnover event occurs when surface waters warm to 4° C. Lakes with this mixing pattern of two stratification periods and two turnover events are called **dimictic** lakes. In shallower lakes, the water column may mix from top to bottom most of the ice-free season with occasional stratification during periods of calm warm conditions. Lakes that mix frequently are termed **polymictic** lakes. In our cold climate, many shallow lakes are **cold monomictic** meaning a thermocline develops every winter, there is one turnover event in spring but the remainder of the ice-free season the lake is polymictic.

Dissolved Oxygen

Oxygen enters a lake at the lake surface and throughout the water column when produced by photosynthesizing plants, including algae, in the lake. Oxygen is consumed within the lake by respiration of living organisms and decomposition of organic material in the lake sediments. In lakes that stratify (see temperature above), oxygen that dissolves into the lake at the surface cannot mix downward into the hypolimnion. At the same time oxygen is depleted in the hypolimnion by decomposition. The result is that the hypolimnion of a lake can become **anoxic**, meaning it contains little or no dissolved oxygen. When a lake is frozen, the entire water column can become anoxic because the surface is sealed off from the atmosphere. Winter anoxic conditions can result in a fish-kill which is particularly common during harsh winters with extended ice-cover. Alberta Surface Water Quality Guidelines suggest dissolved oxygen concentrations (in the epilimnion) must not decline below 5 mg/L and should not average less than 6.5 mg/L over a seven-day period. However, the guidelines also require that dissolved oxygen concentrations remain above 9.5 mg/L in areas where early life stages of aquatic biota, particularly fish, are present.

General Water Chemistry

Water in lakes always contains substances that have been transported by rain and snow or have entered the lake in groundwater and inflow streams. These substances may be dissolved in the water or suspended as particles. Some of these substances are familiar minerals, such as sodium and chloride, which when combined form table salt, but when dissolved in water separate into the two electrically charged components called **ions**. Most dissolved substances in water are in ionic forms and are held in solution due to the polar nature of the water molecule. **Hydrophobic** (water-fearing) compounds such as oils contain little or no ionic character, are non-polar and for this reason do not readily dissolve in water. Although hydrophobic compounds do not readily dissolve, they can still be transported to lakes by flowing water. Within individual lakes, ion concentrations vary from year to year depending on the amount and mineral content of the water entering the lake. This mineral content can be influenced by the amount of precipitation and other climate variables as well as human activities such as fertilizer and road salt application.

Phosphorus and Nitrogen

Phosphorus and nitrogen are important nutrients limiting the growth of algae in Alberta lakes. While nitrogen usually limits agricultural plants, phosphorus is usually in shortest supply in lakes. Even a slight increase of phosphorus in a lake can, given the right conditions, promote algal blooms causing the water to turn green in the summer and impair recreational uses. When pollution originating from livestock manure and human sewage enters lakes not only are the concentrations of phosphorus and nitrogen increased but nitrogen can become a limiting nutrient which is thought to cause blooms of toxic algae belonging to the cyanobacteria. Not all cyanobacteria are toxic, however, the blooms can form decomposing mats that smell and impair dissolved oxygen concentrations in the lake.

Chlorophyll-a

Chlorophyll-*a* is a photosynthetic pigment that green plants, including algae, possess enabling them to convert the sun's energy to living material. Chlorophyll-*a* can be easily extracted from algae in the laboratory. Consequently, chlorophyll-*a* is a good estimate of the amount of algae in the water. Larger aquatic plants, known as macrophytes, rather than algae, dominate some highly productive lakes. In these lakes, chlorophyll-*a* and nutrient values taken from water samples do not include productivity from large aquatic plants. As a result, lakes like Chestermere, which are dominated by macrophytes, can exist at a lower trophic state than if macrophyte biomass was included. Unfortunately, the productivity and nutrient cycling contributions of macrophytes are difficult to sample accurately and are therefore not typically included in trophic state indices.

Secchi Disk Depth

Lakes that are clear are more attractive for recreation, whereas those that are turbid or murky are considered by lake users to have poor water quality. Secchi disk depth is the oldest, simplest, and quickest quantitative measure of water clarity. A Secchi disk is a black and white disk that is lowered down through the water column until it can no longer be seen. Secchi disk depth is the midpoint between the depth at which it disappears when lowered and reappears when it is pulled up again. The Secchi disk depth in lakes with high algal biomass will generally be shallow. However, Secchi disk depth is not only affected by algae. High concentrations of suspended sediments, particularly fine clays or glacial till, are common in plains or mountain reservoirs of Alberta. Mountain reservoirs may have exceedingly shallow Secchi disk depths despite low algal growth and nutrient concentrations.

The euphotic zone, calculated as twice the Secchi disk depth, is the portion of the water column that has sufficient light for aquatic plants to grow. Murky waters, with shallow Secchi depths, can prevent aquatic plants from growing on the lake bottom. Aquatic plants are important because they ensure clear lake water by reducing shoreline erosion and stabilizing lake bottom sediments. Many lakes in Alberta are shallow and have bottom sediments with high concentrations of nutrients. Without aquatic plants, water quality may decline in these lakes due to murky, sediment-laden water and excessive algal blooms. Maintaining aquatic plants in certain areas of a lake is often essential for ensuring good water clarity and a healthy lake as many organisms, like aquatic invertebrates and fish, depend on aquatic plants for food and shelter.

Trophic state

Trophic state is a classification system for lakes that depends on fertility and is a useful index for rating and comparing lakes. From low to high nutrient and algal biomass (as chlorophyll-*a*) concentrations, the trophic states are: **oligotrophic**, **mesotrophic**, **eutrophic** and **hypereutrophic**. The nutrient and algal biomass concentrations that define these categories are shown in table 2 and a graph of Alberta lakes compared by trophic state can be found on the ALMS website. A majority of lakes in Alberta are meso- to eutrophic because they naturally contain high nutrient concentrations due to our deep fertile soils. Thus, lakes in Alberta are susceptible to human impacts because they are already nutrient-rich; any further nutrient increases can bring about undesirable conditions illustrated in Figure 7.




Figure 7: Suggested changes in various lake characteristics with eutrophication. From “Ecological Effects of Wastewater”, 1980.

Table 2: Trophic status based on lake water characteristics

Trophic state	Total Phosphorus (µg/L)	Total Nitrogen (µg/L)	Chlorophyll a (µg/L)	Secchi Depth (m)
Oligotrophic	< 10	< 350	< 3.5	> 4
Mesotrophic	10 - 30	350 - 650	3.5 - 9	4 - 2
Eutrophic	30 - 100	650 - 1200	9 - 25	2 - 1
Hypereutrophic	> 100	> 1200	> 25	< 1

Note: These values are from a detailed study of global lakes reported in Nurnberg 1996. Alberta Environment uses slightly different values for TP and CHL based on those of the OECD reported by Vollenweider (1982). The AENV and OECD cutoffs for TP are 10, 35 and 100; for CHL are 3, 8 and 25. AENV does not have TN or Secchi depth criteria. The corresponding OECD exists for Secchi depth and the cutoffs are 6, 3 and 1.5 m.